



# Knowledge Organisers and Retrieval Questions



Name: \_\_\_\_\_

## Higher Combined Chemistry



# Chapter 1: Atomic structure

## Knowledge organiser


### Development of the model of the atom

#### Dalton's model

John Dalton thought of the \_\_\_\_\_ as a solid sphere that could not be divided into smaller parts. His model did not include \_\_\_\_\_, \_\_\_\_\_, or \_\_\_\_\_.

#### The plum pudding model


Scientists' experiments resulted in the discovery of sub-atomic \_\_\_\_\_ particles. The first to be discovered were electrons – tiny, \_\_\_\_\_ charged particles. The discovery of electrons led to the \_\_\_\_\_ of the atom – a cloud of \_\_\_\_\_ charge, with negative \_\_\_\_\_ embedded in it. \_\_\_\_\_ and \_\_\_\_\_ had not yet been discovered.




#### Alpha scattering experiment

- 1 Scientists fired small, \_\_\_\_\_ charged particles (called \_\_\_\_\_ particles) at a piece of \_\_\_\_\_ foil only a few atoms thick.
- 2 They expected the alpha particles to travel \_\_\_\_\_ through the gold.
- 3 They were surprised that some of the alpha particles \_\_\_\_\_ back and many were \_\_\_\_\_ (alpha scattering).
- 4 To explain why the alpha particles were repelled the scientists suggested that the positive charge and mass of an atom must be concentrated in a small space at its centre. They called this space the \_\_\_\_\_.

#### scientists predicted:



#### actually observed:




#### Nuclear model

Scientists replaced the plum pudding model with the nuclear model and suggested that the electrons \_\_\_\_\_ the nucleus, but not at set \_\_\_\_\_.



#### Electron shell (Bohr) model

Niels Bohr calculated that electrons must orbit the nucleus at fixed distances. These orbits are called \_\_\_\_\_ or \_\_\_\_\_.



#### The proton

Further experiments provided evidence that the nucleus contained smaller particles called \_\_\_\_\_. A proton has an \_\_\_\_\_ charge to an electron.

#### Size

The atom has a radius of \_\_\_\_\_. \_\_\_\_\_ (plural of nucleus) are around \_\_\_\_\_ times smaller than atoms and have a radius of around \_\_\_\_\_.

#### Relative mass

One property of protons, neutrons, and electrons is \_\_\_\_\_ – their masses compared to each other. Protons and neutrons have the same mass, so are given a relative mass of \_\_\_\_\_. It takes almost \_\_\_\_\_ electrons to equal the mass of a single proton – their relative mass is so small that we can consider it as \_\_\_\_\_.

#### The neutron

James Chadwick carried out experiments that gave evidence for a particle with no charge. Scientists called this the \_\_\_\_\_ and concluded that the \_\_\_\_\_ and \_\_\_\_\_ are in the nucleus, and the electrons orbit the nucleus in \_\_\_\_\_.

## Elements and compounds

**Elements** are substances made of one type of atom. Each atom of an element will have the same number of \_\_\_\_\_.

\_\_\_\_\_ are made of different types of atoms chemically \_\_\_\_\_ together. The atoms in a compound have \_\_\_\_\_ numbers of protons.


## Drawing atoms

Electrons in an atom are placed in fixed shells. You can put

- up to \_\_\_\_\_ electrons in the first shell
- eight electrons each in the \_\_\_\_\_ and \_\_\_\_\_ shells.

You must fill up a shell before moving on to the next one.

Name the elements



## Mixtures

A mixture consists of \_\_\_\_\_ elements or compounds that are not \_\_\_\_\_ combined together.

The substances in a mixture can be \_\_\_\_\_ using physical processes.

These processes do not use \_\_\_\_\_ reactions.

## Separating mixtures

- \_\_\_\_\_ – insoluble solids and a liquid
- \_\_\_\_\_ – soluble solid from a solution
- \_\_\_\_\_ – solvent from a solution
- \_\_\_\_\_ – two liquids with similar boiling points
- \_\_\_\_\_ – identify substances from a mixture in solution

## Atoms and particles

	Relative charge	Relative mass	
Proton			= _____ number
Neutron			= _____ number – _____ number
Electron			= same as the number of _____

All atoms have \_\_\_\_\_ numbers of protons and electrons, meaning they have \_\_\_\_\_ overall charge:

*total negative charge from electrons = total positive charge from protons*

## Isotopes

Atoms of the same element can have a different number of \_\_\_\_\_, giving them a different overall \_\_\_\_\_. Atoms of the same element with \_\_\_\_\_ numbers of neutrons are called **isotopes**.

The **relative atomic mass** is the average \_\_\_\_\_ of all the atoms of an element:

$$\text{relative atomic mass} = \frac{(\text{abundance of isotope 1} \times \text{mass of isotope 1}) + (\text{abundance of isotope 2} \times \text{mass of isotope 2})}{100}$$

## Key terms

Make sure you can write a definition for these key terms.

abundance    atom    atomic number    aqueous    compound    electron  
 element    energy level    isotope    neutron    nucleus    orbit  
 product    proton    reactant    relative atomic mass  
 relative charge    relative mass    shell

# Chapter 2: The Periodic Table

## Knowledge organiser

### Development of the Periodic Table

The Periodic Table has changed over time as scientists have organised it differently. Mendeleev was able to accurately predict the properties of undiscovered elements based on the gaps in the table.




	First lists of elements	Mendeleev's Periodic Table	Modern Periodic Table
How are elements ordered?			
Are there gaps?			
How are elements grouped?			
Metals and non-metals			
Problems			

### Group 0

Elements in **Group 0** are called the \_\_\_\_\_. They have the following properties:

- full outer shells with \_\_\_\_\_ electrons, so do not need to \_\_\_\_\_ or \_\_\_\_\_ electrons
- are very \_\_\_\_\_ so exist as single atoms as they do not bond to form molecules
- boiling points that \_\_\_\_\_ down the group.

Name the group 0 elements.



#### Key terms

Make sure you can write a definition for these key terms.

alkali metals    chemical properties    displacement    groups    halogens    inert    isotopes  
noble gas    organised    Periodic Table    reactivity    undiscovered    unreactive

## Group 1 elements

**Group 1** elements react with oxygen, chlorine, and water, for example:

lithium + oxygen  $\rightarrow$  \_\_\_\_\_

lithium + chlorine  $\rightarrow$  \_\_\_\_\_

lithium + water  $\rightarrow$  \_\_\_\_\_ + \_\_\_\_\_

Group 1 elements are called \_\_\_\_\_ because they react with water to form an alkali (a solution of their metal hydroxide).


																H																	He
Li	Be																	B	C	N	O	F	Ne										
Na	Mg																	Al	Si	P	S	Cl	Ar										
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr																
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe																
Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn																
Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg																							

## Group 1 properties

Group 1 elements all have \_\_\_\_\_ electron in their outer shell.

**Reactivity** \_\_\_\_\_ down Group 1 because as you move down the group:

- the atoms \_\_\_\_\_ in size
- the outer electron is further away from the nucleus, and there are more shells \_\_\_\_\_ the outer electron from the nucleus
- the \_\_\_\_\_ attraction between the nucleus and the outer electron is \_\_\_\_\_ so it is easier to lose the one \_\_\_\_\_ electron
- the melting point and boiling point \_\_\_\_\_ down Group 1.



## Group 7 elements


**Group 7** elements are called the \_\_\_\_\_. They are non-metals that exist as molecules made up of \_\_\_\_\_ of atoms.

Name	Formula	State at room temperature	Melting point and boiling point	Reactivity
fluorine				
chlorine				
bromine				
iodine				

## Group 7 reactivity

Reactivity decreases down Group 7 because as you move down the group:

- the atoms \_\_\_\_\_ in size
- the outer shell is further away from the nucleus, and there are more \_\_\_\_\_ between the nucleus and the outer shell
- the electrostatic attraction from the nucleus to the outer shell is \_\_\_\_\_ so it is harder to \_\_\_\_\_ one electron to fill the outer shell.



## Group 7 displacement

More reactive Group 7 elements can take the place of less reactive ones in a compound. This is called displacement.

For example, fluorine displaces chlorine as it is more \_\_\_\_\_:


fluorine + potassium chloride  $\rightarrow$  \_\_\_\_\_ + \_\_\_\_\_

# Chapter 3: Bonding 1

## Knowledge organiser

### Particle model

The three states of matter can be represented in the particle model.



(HT only) This model assumes that:

- 
- 
- 

The amount of energy needed to change the state of a substance depends on the \_\_\_\_\_ between the particles. The stronger the \_\_\_\_\_ between the particles, the \_\_\_\_\_ the melting or boiling point of the substance.

### Covalent bonding

Atoms can \_\_\_\_\_ or \_\_\_\_\_ electrons to form strong chemical bonds.


A **covalent bond** is when electrons are \_\_\_\_\_ between \_\_\_\_\_ atoms.

The number of electrons shared depends on how many \_\_\_\_\_ electrons an atom needs to make a full \_\_\_\_\_.

If you include electrons that are shared between atoms, each atom has a full outer shell.


**Single bond** = each atom shares \_\_\_\_\_ pair of electrons.

**Double bond** = each atom shares \_\_\_\_\_ pairs of electrons.



### Ions

Atoms can \_\_\_\_\_ or \_\_\_\_\_ electrons to give a full outer shell. The number of protons is then \_\_\_\_\_ from the number of electrons. The resulting particle has a charge and is called an \_\_\_\_\_.



### Conductivity

Solid ionic substances do not \_\_\_\_\_ electricity because the ions are fixed in position and not free to carry charge.

When \_\_\_\_\_ or \_\_\_\_\_ in water, ionic substances do \_\_\_\_\_ electricity because the ions are \_\_\_\_\_ to move and carry charge.

### Melting points

Ionic substances have \_\_\_\_\_ melting points because the force of attraction between \_\_\_\_\_ charged ions is strong and so requires lots of energy to break.

### Covalent structures


There are three main types of covalent structure:

#### Structure and bonding

#### Giant covalent

Many billions of atoms, each one with a \_\_\_\_\_ covalent bond to a number of others.


An example of a giant covalent structure is \_\_\_\_\_.



#### Small molecules

Each molecule contains only a few atoms with strong covalent bonds between these atoms. Different molecules are held together by weak \_\_\_\_\_.

For example, \_\_\_\_\_ is made of small molecules.




#### Large molecules

Many repeating units joined by covalent bonds to form a \_\_\_\_\_.

The small section is bonded to many identical sections to the left and right. The 'n' represents \_\_\_\_\_.



Separate chains are held together by intermolecular forces that are stronger than in \_\_\_\_\_ molecules. \_\_\_\_\_ are examples of long molecules.





## Ionic bonding


When metal atoms react with non-metal atoms they \_\_\_\_\_ electrons to the non-metal atom.



Metal atoms \_\_\_\_\_ electrons to become \_\_\_\_\_ ions.  
Non-metal atoms \_\_\_\_\_ electrons to become \_\_\_\_\_ ions.

## Giant ionic lattice

When metal atoms \_\_\_\_\_ electrons to non-metal atoms you end up with \_\_\_\_\_ and \_\_\_\_\_ ions. These are attracted to each other by the strong **electrostatic force of attraction**. This is called \_\_\_\_\_.




The electrostatic force of attraction works in all \_\_\_\_\_, so many billions of ions can be bonded together in a \_\_\_\_\_ structure.

## Formulae

The formula of an ionic substance can be worked out

- from its \_\_\_\_\_:  
for every \_\_\_\_\_ magnesium ion there are \_\_\_\_\_ fluoride ions – so the formula for magnesium fluoride is  $\text{MgF}_2$
- from a \_\_\_\_\_:  
there are nine  $\text{Fe}^{2+}$  ions and 18  $\text{S}^{2-}$  ions – simplifying this ratio gives a formula of \_\_\_\_\_




## Metals: Structure and properties

The atoms that make up metals form \_\_\_\_\_. The electrons in the outer shells of the atoms are \_\_\_\_\_ – this means they are free to move through the \_\_\_\_\_ structure.

The positive metal ions are then attracted to these delocalised electrons by the \_\_\_\_\_.

Some important properties of metals are:

- 
- 
- 



# Chapter 3: Bonding 2

## Knowledge organiser


Properties	High melting and boiling points because the _____ covalent bonds between the atoms must be _____ to melt or boil the substances.  This requires a lot of _____. _____ at room temperature.	_____ melting and boiling points because only the _____ forces need to be overcome to melt or boil the substances, not the _____ between the atoms.  This does not require a lot of energy as the intermolecular forces are _____.  Normally _____ or _____ at room temperature.	Melting and boiling points are _____ compared to giant _____ substances but higher than for small molecules.  Large molecules have _____ intermolecular forces than small molecules, which require more _____ to overcome.  Normally _____ at room temperature.
	Most covalent structures do not _____ electricity because they do not have _____ or _____ that are free to move to carry charge.		

### Graphite

Graphite is a giant covalent structure, but is different to other giant covalent substances.

#### Structure

Made only of \_\_\_\_\_ – each carbon atom bonds to \_\_\_\_\_ others, and forms hexagonal rings in \_\_\_\_\_. Each carbon atom has one \_\_\_\_\_ electron, which is delocalised and therefore \_\_\_\_\_ to move around the structure.



#### Hardness

The layers can \_\_\_\_\_ over each other because they are not \_\_\_\_\_ bonded. Graphite is therefore softer than \_\_\_\_\_, even though both are made only of carbon, as each atom in diamond has \_\_\_\_\_ strong covalent bonds.

#### Conductivity

The \_\_\_\_\_ electrons are free to move through graphite, so can carry charges and allow an electrical current to \_\_\_\_\_. Graphite is therefore a \_\_\_\_\_ of electricity.

### Graphene

Graphene consists of only a single layer of \_\_\_\_\_. Its strong covalent bonds make it a \_\_\_\_\_ material that can also conduct electricity. It could be used in composites and high-tech electronics.

### Fullerenes


- \_\_\_\_\_ cages of carbon atoms bonded together in one molecule
- can be arranged as a \_\_\_\_\_ or a \_\_\_\_\_ (called a **nanotube**)
- molecules held together by \_\_\_\_\_ forces, so can slide over each other
- conduct electricity

#### Spheres


Buckminsterfullerene was the first fullerene to be discovered, and has \_\_\_\_\_ carbon atoms.

Other fullerenes exist with different numbers of carbon atoms arranged in rings that form hollow shapes.

Fullerenes like this can be used as \_\_\_\_\_ and in \_\_\_\_\_.



#### Nanotubes




The carbon atoms in nanotubes are arranged in cylindrical tubes.

Their high \_\_\_\_\_ strength (they are difficult to break when pulled) makes them useful in electronics.



## Alloys

Pure metals are often too \_\_\_\_\_ to use as they are. Adding atoms of a different element can make the resulting mixture \_\_\_\_\_ because the new atoms will have a different \_\_\_\_\_ to the pure metal's atoms. This will disturb the \_\_\_\_\_ arrangement of the layers, preventing them from \_\_\_\_\_ over each other. The harder mixture is called an **alloy**.



## Measuring particles

We use different units and scales to measure the size of particles.

Particle	Particulate matter	Size	Standard form	Full form
grain of sand	N/A	0.1 mm	$1 \times 10^{-4}$ m	0.0001 m
coarse particles (e.g. dust)	PM <sub>10</sub>	10 $\mu$ m	$1 \times 10^{-5}$ m	0.00001 m
fine particles	PM <sub>2.5</sub>	100 nm	$1 \times 10^{-7}$ m	0.0000001 m
nanoparticles	< PM <sub>2.5</sub>	1 to 100 nm	$1 \times 10^{-9}$ to $1 \times 10^{-7}$ m	0.000000001 m to 0.0000001 m

PM stands for \_\_\_\_\_ and is another way of measuring very small particles.

## Uses of nanoparticles

Nanoparticles often have very different \_\_\_\_\_ to bulk materials of the same substance, caused by their high surface area-to-volume ratio.

Nanoparticles have many uses and are an important area of research. They are used in \_\_\_\_\_, \_\_\_\_\_, and as \_\_\_\_\_.

However, nanoparticles have the potential to be hazardous to health and to \_\_\_\_\_, so it is important that they are researched further.

## Key terms

Make sure you can write a definition for these key terms.

- conductivity
- conductor
- delocalised electron
- electrostatic force of attraction
- ion
- lattice
- layer
- malleable
- nanoparticle
- particulate matter
- surface area to volume ratio
- transfer

# Chapter 4: Calculations

## Knowledge organiser

### Formula mass

Every substance has a \_\_\_\_\_,  $M_r$ .

$M_r =$  \_\_\_\_\_

### Avogadro's constant (HT only)

One mole of a substance contains \_\_\_\_\_ atoms, ions, or molecules.  
This is **Avogadro's constant**.

One mole of a substance has the same \_\_\_\_\_ as the  $M_r$  of the substance.  
For example, the  $M_r$  ( $H_2O$ ) = \_\_\_\_\_, so \_\_\_\_\_ g of water molecules contains  
\_\_\_\_\_ molecules, and is called one \_\_\_\_\_ of water.

You can write this as: moles = \_\_\_\_\_

### Using balanced equations (HT)

In a balanced symbol equation the sum of the

Write down the steps for how to calculate the mass of a product from a given mass of a specific reactant.

1. \_\_\_\_\_
2. \_\_\_\_\_
3. \_\_\_\_\_
4. \_\_\_\_\_

### Theoretical yield

The **theoretical yield** of a chemical reaction is the mass of a product that you expect to be \_\_\_\_\_.

Even though no \_\_\_\_\_ are gained or lost during a chemical reaction, it is not always possible to obtain the theoretical yield because:

### Percentage yield

The **yield** is the amount of \_\_\_\_\_ that you actually get in a chemical reaction.

Percentage yield is the actual yield as a proportion of the theoretical yield:

percentage yield = \_\_\_\_\_

### Atom economy

The **atom economy** of a reaction tells you the \_\_\_\_\_ of atoms that you started with that are part of \_\_\_\_\_ products.

High atom economies are more sustainable, as they mean \_\_\_\_\_ atoms are being wasted in products that are not useful.

The percentage atom economy is calculated by:

atom economy = \_\_\_\_\_

### Concentration

Concentration is the amount of solute in a \_\_\_\_\_ of solvent.

The unit of concentration is  $g/dm^3$ .

Concentration can be calculated using:

concentration ( $g/dm^3$ ) = \_\_\_\_\_

Sometimes volume is measured in  $cm^3$ :

volume ( $dm^3$ ) = \_\_\_\_\_

- lots of solute in little solution = \_\_\_\_\_
- little solute in lots of solution = \_\_\_\_\_

### Moles of gases (HT only)

At any given temperature and pressure, the same number of moles of a gas will \_\_\_\_\_ the same volume.

At room temperature (\_\_\_\_\_) and pressure (\_\_\_\_\_), one mole of any gas will occupy \_\_\_\_\_.

To calculate the number of moles of a gas:

moles of a gas = \_\_\_\_\_  
or  $24 dm^3$

moles of a gas = \_\_\_\_\_  
 $cm^3$



### Key terms

Make sure you can write a definition for these key terms.

atom economy    burette    concordant    end point  
excess reactant    formula mass    limiting reactant  
percentage yield    pipette    room temperature and pressure  
theoretical yield    titration    titre    useful    yield

ly)

of the reactants equals the sum of the  $M_r$  of the products.

Write down the steps for how to balance an equation.

1. \_\_\_\_\_
2. \_\_\_\_\_
3. \_\_\_\_\_
4. \_\_\_\_\_

## Excess and limiting reactants (HT only)

In a chemical reaction between two or more reactants, often one of the reactants will run out before the others. You then have some of the other reactants left over. The reactant that is left over is in \_\_\_\_\_. The reactant that runs out is the \_\_\_\_\_.

Write down the steps for working out which reactants are in excess, and which is the limiting reactant.

1. \_\_\_\_\_
2. \_\_\_\_\_
3. \_\_\_\_\_
4. \_\_\_\_\_
5. \_\_\_\_\_

## Concentration in $\text{mol/dm}^3$

Concentration can also be measured in  $\text{mol/dm}^3$ .

concentration of solution ( $\text{mol/dm}^3$ ) =

You can use this formula and mass = moles  $\times M_r$  to calculate

- The greater the mass of solute in solution, the greater the number of \_\_\_\_\_ of solute, and therefore the greater the \_\_\_\_\_.
- If the \_\_\_\_\_ number moles of solute is dissolved in a smaller volume of solution, the concentration will be \_\_\_\_\_.

mol is a  
the unit  
of moles

## Calculating concentration

To calculate the concentration of the unknown solution (the solution in the conical flask):

- 1 Write a \_\_\_\_\_ for the reaction.
- 2 Calculate the moles used from the known solution using:

moles =


- 3 Use the \_\_\_\_\_ from the balanced symbol equation to deduce the number of moles present in the unknown solution.
- 4 Calculate the concentration of the unknown solution using:

concentration ( $\text{mol/dm}^3$ ) =

## Titration

**Titration** is an experimental technique to work out the concentration of an \_\_\_\_\_ solution in the reaction between an \_\_\_\_\_ and an \_\_\_\_\_.

- 1 Use a \_\_\_\_\_ to extract a known volume of the solution with an unknown concentration. A pipette measures a \_\_\_\_\_ volume only.
- 2 Add the solution of unknown concentration to a \_\_\_\_\_ flask and put the conical flask on a white tile.
- 3 Add a few \_\_\_\_\_ of a suitable indicator to the conical flask.
- 4 Add the other solution with a known concentration to the \_\_\_\_\_.
- 5 Carry out a rough titration to find out approximately what volume of solution in the burette needs to be added to the solution in the conical flask. Add the solution from the burette to the solution in the conical flask  $1\text{cm}^3$  at a time until the \_\_\_\_\_ is reached.
- 6 The \_\_\_\_\_ is when the indicator just changes \_\_\_\_\_.
- 7 Record the \_\_\_\_\_ of the end point as your rough value.
- 8 Now repeat steps 1-7, but as you approach the end point add the solution from the burette drop-by-drop. \_\_\_\_\_ the conical flask in between drops.
- 9 Record the volume of the end point.



# Chapter 5: Chemical changes 1


## Knowledge organiser

### Reactions of metals

The \_\_\_\_\_ of a metal is how chemically reactive it is. When added to water, some metals react very vigorously – these metals have \_\_\_\_\_ reactivity. Other metals will barely react with water or acid, or won't react at all – these metals have \_\_\_\_\_ reactivity.

### Reactivity series

The reactivity series places \_\_\_\_\_ in order of their reactivity. Sometimes, for example in the table below, hydrogen and carbon are included in the series, even though they are \_\_\_\_\_.


Reaction with water	Reaction with acid	Reactivity series		Extraction method
		Metal	Reactivity	
		potassium		
		sodium		
		lithium		
		calcium		
		magnesium		
		aluminium (carbon) zinc		
		iron		
		tin		
		lead (hydrogen) copper		
		silver		
		gold		

### Metal extraction

Some metals, like gold, are so \_\_\_\_\_ that they are found as pure metals in the Earth's \_\_\_\_\_ and can be mined.

Most metals exist as compounds in rock and have to be \_\_\_\_\_ from the rock. If there is enough metal compound in the rock to be worth extracting it is called an \_\_\_\_\_.

Metals that are \_\_\_\_\_ reactive than carbon can be extracted by \_\_\_\_\_ with carbon. For example:




Metals that are more reactive than carbon can be extracted using a process called \_\_\_\_\_.

### Reduction and oxidation


If a substance gains oxygen in a reaction, it has been \_\_\_\_\_.

If a substance loses oxygen in a reaction, it has been \_\_\_\_\_.

For example:



iron has been \_\_\_\_\_



\_\_\_\_\_ has been reduced

### Salts

When acids react with metals or metal compounds, they form \_\_\_\_\_. A salt is a compound where the \_\_\_\_\_ from an acid has been replaced by a metal. For example nitric acid,  $\text{HNO}_3$ , reacts with sodium to form \_\_\_\_\_. The H in nitric acid is replaced with Na.


The table shows how to name salts.

Acid	hydrochloric acid	sulfuric acid	nitric acid
Formula			
Ions formed in solution			
Type of salt formed			
Sodium salt example			

## Displacement reactions

In a **displacement** reaction a \_\_\_\_\_ reactive element takes the place of a *less* reactive element in a compound.

For example:



\_\_\_\_\_ is more reactive than copper, so iron displaces the copper in copper sulfate.

## Ionic equations (HT only)

When an ionic compound is dissolved in a solution, we can write the compound as its separate ions. For example,  $\text{CuSO}_4(\text{aq})$  can be written as \_\_\_\_\_ and \_\_\_\_\_.

The displacement reaction of copper sulfate and iron can be written as:

The  $\text{SO}_4^{2-}$  is unchanged in the reaction – it is a \_\_\_\_\_. Spectator ions are removed from the equation to give an **ionic equation**:

Metals, covalent substances, and solid ionic substances do not split into ions in the ionic equation.

## Half equations (HT only)

In the displacement reaction, an iron atom loses two electrons to form a iron ion:



A copper ion gains two electrons to form a copper atom:



These two equations are called \_\_\_\_\_ – they each show half of the ionic equation.

## Reactivity and ions

A metal's reactivity depends on how readily it forms an \_\_\_\_\_ by losing electrons.

In the displacement reaction of copper sulfate and iron, iron forms an ion more easily than copper.

At the end of the reaction you are left with \_\_\_\_\_ ions, not \_\_\_\_\_ ions.

## Steps for writing an ionic equation (HT only)

- 1.
- 2.
- 3.
- 4.

## Reduction and oxidation: electrons (HT only)

Oxidation and reduction (**redox** reactions) can be defined in terms of oxygen, but can also be defined as the \_\_\_\_\_ or \_\_\_\_\_ of electrons.

Oxidation is the \_\_\_\_\_ of electrons, and reduction is the \_\_\_\_\_ of electrons.

In the example displacement reaction:

- iron atoms have been \_\_\_\_\_
- copper ions have been \_\_\_\_\_.

## Acids and alkalis

**Acids** are compounds that, when dissolved in water, release \_\_\_\_\_ ions. There are three main acids: \_\_\_\_\_, \_\_\_\_\_, and \_\_\_\_\_.

**Alkalis** are compounds that, when dissolved in water, release \_\_\_\_\_ ions.

The **pH** scale is a measure of acidity and alkalinity. It runs from 1 to 14.


- Aqueous solutions with  $\text{pH} < 7$  are \_\_\_\_\_.
- Aqueous solutions with  $\text{pH} > 7$  are \_\_\_\_\_.
- Aqueous solutions with  $\text{pH} = 7$  are \_\_\_\_\_.

## Indicators

Indicators can show if something is an acid or an alkali.

- \_\_\_\_\_ can also tell us the approximate pH of a solution.
- \_\_\_\_\_ can give us the exact pH of a solution.

### The pH scale



# Chapter 5: Chemical changes 2

## Knowledge organiser

### Reactions of acids

#### Reactions of acids with metals

Acids react with some metals to form \_\_\_\_\_ and \_\_\_\_\_.

magnesium + hydrochloric acid → \_\_\_\_\_

#### Neutralisation reactions

##### Reactions of acids with metal hydroxides

Acids react with metal hydroxides to form \_\_\_\_\_ and \_\_\_\_\_.

hydrochloric acid + sodium hydroxide → \_\_\_\_\_

The ionic equation for this reaction is always:

\_\_\_\_\_

##### Reactions of acids with metal oxides

Acids react with metal oxides to form \_\_\_\_\_ and \_\_\_\_\_.

\_\_\_\_\_ + \_\_\_\_\_ → \_\_\_\_\_ + \_\_\_\_\_

##### Reactions of acids with metal carbonates

Acids react with metal carbonates to form a \_\_\_\_\_, \_\_\_\_\_ and \_\_\_\_\_.

hydrochloric acid + sodium carbonate → \_\_\_\_\_


### Alkalis and bases

\_\_\_\_\_ neutralise acids to form water in **neutralisation** reactions. Some metal hydroxides dissolve in water to form \_\_\_\_\_ solutions, called alkalis.

Some metal oxides and metal hydroxide do not dissolve in \_\_\_\_\_. They are **bases**, but are not alkalis.



### Balancing symbol equations

When writing symbol equations you need to ensure that the number of each \_\_\_\_\_ on each side is \_\_\_\_\_.




unbalanced

there are \_\_\_\_\_ hydrogen atoms on each side, but \_\_\_\_\_ oxygen atoms in the reactants and \_\_\_\_\_ in the product



balanced

there are \_\_\_\_\_ hydrogen atoms on each side, and \_\_\_\_\_ oxygen atoms on each side



### State symbols

A balanced symbol equation should also include state symbols. Complete the table.

State	Symbol



#### Key terms

Make sure you can write a definition for these key terms.

displacement

electrolysis

extraction

half equation

ion

ionic equation

metal ore

oxidation

reactivity

reactivity series

redox

reduction

spectator ion

state symbols





**strong and weak acids**

Sulfuric acid, nitric acid, and hydrochloric acid, are all **strong acids**. This means that, when dissolved in water, every molecule splits up into ions – they are completely ionised:

$\text{H}_2\text{SO}_4(\text{aq}) \rightarrow 2\text{H}^+(\text{aq}) + \text{SO}_4^{2-}(\text{aq})$

$\text{HNO}_3(\text{aq}) \rightarrow \text{H}^+(\text{aq}) + \text{NO}_3^-(\text{aq})$

$\text{HCl}(\text{aq}) \rightarrow \text{H}^+(\text{aq}) + \text{Cl}^-(\text{aq})$

Weak acids, such as ethanoic acid, are **weak acids**. This means that only a percentage of their molecules split up into ions when dissolved in water – they are **partially ionised**.


For a given concentration, the **stronger** the acid, the **lower** the pH.

**concentrated and dilute acids**

**Concentration** tells us how much of a substance there is dissolved in water:

more concentrated acids have lots of acid in a **small** volume of water

less concentrated acids (dilute acids) have little acid in a large volume of water.



**crystallisation**



You can produce a solid salt from an insoluble base by **crystallisation**.

The experimental method is:

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_


\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_




# Chapter 6: Electrolysis

## Knowledge organiser

### Electrolysis

In the process of **electrolysis**, an electric \_\_\_\_\_ is passed through an **electrolyte**. An electrolyte is a \_\_\_\_\_ that contains \_\_\_\_\_ and so can conduct electricity. This causes the ions to move to the \_\_\_\_\_, where they form pure elements.

Label the diagram.



### Electrolysis of molten compounds

Solid ionic compounds do not conduct electricity as the ions cannot move. To undergo electrolysis they must be \_\_\_\_\_ or dissolved, so the ions are free to move.

When an ionic compound is molten:

- The positive metal ions are *attracted* to the \_\_\_\_\_, where they will \_\_\_\_\_ electrons to form the pure metal
- The negative non-metal ions are *attracted* to the \_\_\_\_\_, where they will \_\_\_\_\_ electrons and become the pure non-metal.

For example, molten sodium chloride, NaCl, can undergo electrolysis to form \_\_\_\_\_ at the cathode and \_\_\_\_\_ at the anode.

### Half equations (HT only)

sodium chloride → sodium + chlorine

\_\_\_\_\_ → \_\_\_\_\_ + \_\_\_\_\_

- at the cathode:

\_\_\_\_\_ + \_\_\_\_\_ → \_\_\_\_\_


- at the anode:

\_\_\_\_\_ → \_\_\_\_\_ + \_\_\_\_\_

### Electrolysis of aqueous solutions

Solid ionic compounds can also undergo electrolysis when dissolved in water.

- It requires \_\_\_\_\_ energy to dissolve ionic compounds in water than it does to melt them.
- However, in the electrolysis of solutions, the pure elements are not always produced. This is because the water can also undergo \_\_\_\_\_:



### Products at the anode

In the electrolysis of a solution, if the non-metal contains oxygen then oxygen \_\_\_\_\_ is formed at the anode:

- The \_\_\_\_\_ ions formed from the ionisation of water are attracted to the anode.
- The  $\text{OH}^-(\text{aq})$  ions \_\_\_\_\_ electrons to the anode and form oxygen gas.
- $4\text{OH}^-(\text{aq}) \rightarrow$  \_\_\_\_\_


If the non-metal ion is a halogen, then the halogen gas is formed at the anode.

- $2\text{Cl}^-(\text{aq}) \rightarrow$  \_\_\_\_\_

### Products at the cathode

In the electrolysis of a solution, if the metal is \_\_\_\_\_ **reactive** than hydrogen then hydrogen gas is formed at the cathode:

- The \_\_\_\_\_ ions from the ionisation of water are attracted to the cathode and react with it.
- The  $\text{H}^+(\text{aq})$  ions \_\_\_\_\_ electrons from the cathode and form hydrogen gas.
- $2\text{H}^+(\text{aq}) +$  \_\_\_\_\_
- The metal ions remain in \_\_\_\_\_.



## Electrolysis of aluminium oxide

Electrolysis can be used to extract metals from their ionic compounds.

Electrolysis is used if the metal is more reactive than \_\_\_\_\_.


Aluminium is extracted from \_\_\_\_\_ by electrolysis.

- 1 The aluminium oxide is mixed with a substance called \_\_\_\_\_, which lowers the melting point.
- 2 The mixture is then heated until it is \_\_\_\_\_.
- 3 The resulting molten mixture undergoes electrolysis.


aluminium oxide  $\rightarrow$  aluminium + oxygen



cathode: pure aluminium is formed




anode: oxygen is formed



In the electrolysis of aluminium, the anode is made of \_\_\_\_\_.


The graphite reacts with the \_\_\_\_\_ to form \_\_\_\_\_ and so slowly wears away. It therefore needs to be replaced frequently.



## Electrolysis of zinc chloride

Molten zinc chloride is broken down by electrolysis. This means

\_\_\_\_\_ metal is collected at the \_\_\_\_\_ and a \_\_\_\_\_ is collected at the \_\_\_\_\_. Free \_\_\_\_\_ from the molten zinc chloride are able to move around and carry \_\_\_\_\_, hence why the bulb lights up. Label the diagram.



### Key terms

Make sure you can write a definition for these key terms.

anode

cathode

cryolite

electrode

electrolysis

electrolyte

reactivity

# Chapter 7: Energy changes

## Knowledge organiser

### Energy changes

During a chemical reaction, energy transfers occur.

Energy can be transferred:

- to the surroundings – \_\_\_\_\_
- from the surroundings – \_\_\_\_\_

This energy transfer can cause a \_\_\_\_\_ change.

Energy is always conserved in chemical reactions.

This means that there is the \_\_\_\_\_

### The surroundings

When chemists say energy is transferred from or to “the surroundings” they mean \_\_\_\_\_.


For example, imagine you have a reaction mixture in a test tube. If you measure the temperature in the test tube using a thermometer, the thermometer is then part of the surroundings.

- If the thermometer records an increase in temperature, the reaction in the test tube is \_\_\_\_\_.
- If the thermometer records a decrease in temperature, the reaction in the test tube is \_\_\_\_\_.

### Reaction profiles

A \_\_\_\_\_ shows whether a reaction is exothermic or endothermic.

The \_\_\_\_\_ is the minimum amount of energy that particles must have to react when they \_\_\_\_\_.



### Bonds (HT only)

Atoms are held together by strong \_\_\_\_\_ bonds. In a reaction, those bonds are \_\_\_\_\_ and new ones are made between \_\_\_\_\_ atoms.

- Breaking a bond \_\_\_\_\_ energy so is \_\_\_\_\_.
- Making a bond \_\_\_\_\_ energy so is \_\_\_\_\_.

### Breaking bonds

If a lot of energy is released when \_\_\_\_\_ the bonds and only a \_\_\_\_\_ energy is required to break them, then overall energy is released and the reaction as a whole is \_\_\_\_\_.

### Making bonds

If a \_\_\_\_\_ energy is released when making the bonds and a \_\_\_\_\_ is required to break them, then overall energy is taken in and the reaction as a whole is \_\_\_\_\_.

### Bond calculations

Different bonds require different amounts of energy to be broken (their **bond energies**). To work out the overall energy change of a reaction, you need to:

- 1 work out how much energy is required to break all the bonds in the \_\_\_\_\_
- 2 work out how much energy is \_\_\_\_\_ when making all the bonds in the products.

overall energy transferred = \_\_\_\_\_ - \_\_\_\_\_

- A \_\_\_\_\_ number means an endothermic reaction.
- A \_\_\_\_\_ number means an exothermic number.

Reaction	Energy transfer	Temperature change	Example	Everyday use	Bonds
exothermic			• • •	• •	
endothermic			• •	•	

## Chemical cells

In a metal displacement reaction, one metal is \_\_\_\_\_. These electrons are \_\_\_\_\_ to another metal, which gains the electrons and so is \_\_\_\_\_.


By using a **chemical cell** to conduct this reaction, the electron's movement generates a \_\_\_\_\_.

In the cell shown, the zinc atoms from the electrode lose electrons, turn into \_\_\_\_\_, and move into the solution.

The \_\_\_\_\_ travel through the circuit to the \_\_\_\_\_ electrode, causing the LED to \_\_\_\_\_.

Once at the copper electrode, a metal ion from the solution will pick the electrons up and become a \_\_\_\_\_ atom.

The greater the difference in reactivity between the two metals in the cell, the greater the \_\_\_\_\_ produced.



## Batteries


A **battery** is formed of \_\_\_\_\_ cells connected in \_\_\_\_\_.

- Some batteries are \_\_\_\_\_. An external electric current is applied, which reverses the reaction.
- Some batteries, like \_\_\_\_\_ batteries, are not rechargeable because the reaction is not \_\_\_\_\_. Once the reactants are used up, the chemical reaction \_\_\_\_\_ and no more potential differences are \_\_\_\_\_.

## Hydrogen fuel cells


**Fuel cells** use a \_\_\_\_\_ and \_\_\_\_\_ from the air to generate a potential difference.

\_\_\_\_\_ fuel cells generate electricity from hydrogen and oxygen. The overall reaction is:



The hydrogen is oxidised to produce \_\_\_\_\_.

There are different types of hydrogen fuel cell. In alkaline fuel cells, the half equations are below:



### Advantages

• \_\_\_\_\_  
• \_\_\_\_\_  
• \_\_\_\_\_

### Disadvantages

• \_\_\_\_\_  
• \_\_\_\_\_  
• \_\_\_\_\_



### Key terms

**Make sure you can write a definition for these key terms.**

activation energy    battery  
bond energy    chemical cell  
combustion    endothermic  
exothermic    fuel cell  
neutralisation    oxidation  
reaction profile    rechargeable  
thermal decomposition

# Chapter 8: Rates and equilibrium 1

## Knowledge organiser

### Rates of reaction

The **rate of a reaction** is how quickly the \_\_\_\_\_ turn into the \_\_\_\_\_.

To calculate the rate of a reaction, you can measure:

- how quickly a reactant is used up

mean rate of reaction = \_\_\_\_\_

- how quickly a product is produced.

mean rate of reaction = \_\_\_\_\_

For reactions that involve a gas, this can be done by \_\_\_\_\_


### Volume of gas produced

The reaction mixture is connected to a \_\_\_\_\_ or an upside down measuring cylinder. As the reaction proceeds the gas is collected.

The rate for the reaction is then:


rate = \_\_\_\_\_

Volume is measured in \_\_\_\_\_ and time in \_\_\_\_\_, so the unit for rate is \_\_\_\_\_.



### Change in mass

The reaction mixture is placed on a mass balance. As the reaction proceeds and the \_\_\_\_\_ product is given off, the mass of the flask will \_\_\_\_\_.



The rate for the reaction is then:

rate = \_\_\_\_\_

The mass is measured in grams and time is measured in seconds. Therefore, the unit of rate is \_\_\_\_\_.

### Calculating rate from graphs (HT only)

The results from an experiment can be plotted on a graph.



- A steep gradient means \_\_\_\_\_
- A shallow gradient means \_\_\_\_\_

### Mean rate at specific time

To obtain the rate at a specific time draw a \_\_\_\_\_ to the graph and calculate its \_\_\_\_\_.

### Mean rate between two points in time

To get the mean rate of reaction between two points in time:



Rate at 50 s = \_\_\_\_\_ = \_\_\_\_\_

The gradient is the change in y divided by the change in x for a right-angled triangle drawn from the tangent.



## Collision theory

For a reaction to occur, the reactant particles need to \_\_\_\_\_. When the particles collide, they need to have enough \_\_\_\_\_ to react or they will just bounce apart. This amount of energy is called the \_\_\_\_\_.

You can increase the rate of a reaction by:

- \_\_\_\_\_
- \_\_\_\_\_
- \_\_\_\_\_



## Factors affecting rate of reaction


Condition that increases rate	How is this condition caused?	Why it has that effect
increasing the temperature		1 2
increasing the concentration of solutions		
increasing the pressure of gases		
increasing the surface area of solids		



## Catalysts

Some reactions have specific substances called \_\_\_\_\_ that can be added to increase the rate. These substances are not used up in the reaction.

A catalyst provides a different \_\_\_\_\_ that has a \_\_\_\_\_ activation energy. As such, more particles will collide with enough energy to react, so more collisions result in a reaction.



# Chapter 8: Rates and equilibrium 2

## Knowledge organiser

### Reaction conditions

The conditions of a reaction refer to the external environment of the reaction. When the reaction occurs in a closed system, you can change the conditions by:

- 
- 
- 

### Le Châtelier's principle (HT only)

At equilibrium, the amount of reactants and products is constant. In order to change the amounts of reactant and product at equilibrium the \_\_\_\_\_ of the reaction must be changed. The closed system will then counteract the change by favouring either the \_\_\_\_\_ or the \_\_\_\_\_. This is known as **Le Châtelier's principle**.

### Changing concentrations (HT only)

Change	Effect	Explanation
decrease concentration of product		
increase concentration of product		
decrease concentration of reactant		
increase concentration of reactant		

### Changing temperature (HT only)

Change	Effect	Explanation
increase temperature of surroundings		
decrease temperature of surroundings		

### Changing pressure (HT only)

Change	Effect	Explanation
increase the pressure		
decrease the pressure		




#### Key terms

Make sure you can write a definition for these key terms.

activation energy   catalyst   collision   collision theory   closed system  
conditions   dynamic equilibrium   frequency of collision   gradient  
Le Châtelier's principle   rate of reaction   reversible reaction   tangent

# reversible reactions

some reactions, the products can react to produce the original reactants. This is called a **reversible reaction**. When writing chemical equations for reversible reactions, use the  $\rightleftharpoons$  symbol.



this reaction:  
 A and B can react to form C and D – the \_\_\_\_\_ reaction.  
 C and D can react to form A and B – the \_\_\_\_\_ reaction.  
 The different directions of the reaction have \_\_\_\_\_ energy changes.  
 The forward reaction is \_\_\_\_\_, the reverse reaction will be *exothermic*.  
 The same amount of energy is transferred in each \_\_\_\_\_.

# Equilibrium

In a \_\_\_\_\_ system no reactants or products can escape. If a reversible reaction is carried out in a closed system, it will eventually reach \_\_\_\_\_ – a point in time when the forward and reverse reactions have the same rate.

At dynamic equilibrium:

- 
- 
- 

# Dynamic equilibrium

At dynamic equilibrium the amount of reactant and product are constant, but not necessarily \_\_\_\_\_.

You could have a mixture of reactants and products in a 50:50 ratio, in a 75:25 ratio, or in any ratio at all. The \_\_\_\_\_ of the reaction are what change that ratio.

# how dynamic equilibrium is reached

Progress of reaction	start of reaction	middle of reaction	at dynamic equilibrium
Amount of A + B	high	decreasing	_____
frequency of collisions A + B	high	_____	_____
Rate of forward reaction	_____	_____	_____
	<div> </div>		
Amount of C + D	_____	_____	_____
frequency of collisions C + D	_____	_____	constant
Rate of reverse reaction	_____	_____	_____

# Chapter 9: Crude oils and fuels

## Knowledge organiser

### Crude oil

**Crude oil** is incredibly important to our society and economy. It is formed from the remains of ancient \_\_\_\_\_ - living organisms (mostly \_\_\_\_\_) that died \_\_\_\_\_ of years ago.

Raw crude oil is a thick black \_\_\_\_\_ made of a large number of different \_\_\_\_\_ mixed together. Most of the compounds are \_\_\_\_\_ of various sizes. Hydrocarbons are \_\_\_\_\_.

### Combustion

Hydrocarbons are used as \_\_\_\_\_. This is because when they react with \_\_\_\_\_ they release a lot of \_\_\_\_\_. This reaction is called \_\_\_\_\_. Complete combustion is a type of combustion where the only products are \_\_\_\_\_ and \_\_\_\_\_.

### Properties

Whether or not a particular hydrocarbon is useful as a fuel depends on its properties:





- \_\_\_\_\_ - how easily it burns
- \_\_\_\_\_ - the temperature at which it boils
- \_\_\_\_\_ - how thick it is

Its properties in turn depend on the length of the molecule. Complete the table.

Chain length	Flammability	Boiling point	Viscosity
long chain	low		
short chain			

### Alkanes

One \_\_\_\_\_ of hydrocarbon molecules are called **alkanes**. Alkane molecules only have \_\_\_\_\_ bonds in them. The first four alkanes are:



The different alkanes have different numbers of carbon atoms and hydrogen atoms. You can always work the molecular formula of an alkane by using \_\_\_\_\_.



#### Key terms

Make sure you can write a definition for these key terms.

alkanes    alkenes    boiling point    combustion    cracking    crude oil    feedstock  
flammability    fractional distillation    fuel    hydrocarbon    viscosity